

Jeffrey H. Berg, M.D. Stephanie Clop, M.D. Thomas B. Fleeter, M.D. George Kartalian, M.D. David R. Miller, M.D. Dhruv B. Pateder, M.D. James D. Reeves, MD Raymond Thal, M.D.

1800 Town Center Drive, Suite 300 Reston VA 20190 • 6201 Centreville Road, Suite 600 Centreville VA 20121 Phone: 703-435-6604 • Fax: 703-787-6575 Phone: 703- 378- 4860 • Fax: 703-378-4868

Bankart Repair Rehabilitation

I. Week 1

- Wear sling day and night
- Post-op PT appointment at 2nd post-op day
- Cervical stretches
- Active elbow flexion and wrist/hand ROM
- Begin grip and wrist strengthening exercises
- Cryotherapy (GAME READY) and Electrical stimulation for edema and pain control prin
- Pendulums and continue with HEP

II. 3 Weeks from Surgery

- Resume physical therapy sessions 1-3 x/week
- Isometric shoulder pinches/retraction
- Focus PROM and AAROM to 90 degrees for flexion and ER at neutral to 30 degrees, abduction to 60 degrees
- Pulleys for AAROM abduction, scaption and flexion pain free range (issue pulleys for HEP prn)
- Seated unilateral scapular retraction in gravity eliminated position (i.e. arm resting on table)
- Bicep/Triceps isometric strengthening multi-angle
- Sidelying scapular mobilizations

III. 4 Weeks from Surgery

- Neutral shoulder isometrics 6 directions
- Supine wand exercises flexion to 90 degrees, ER at 0 degrees of abduction to 30 degrees, abduction to 60 degrees
- Discontinue sling use at home and sleeping but continue wear public places
- Sidelying scapular PNF (anterior elevation/posterior depression & posterior elevation /anterior depression)
- AAROM with wall walking flexion and abduction
- Progress passive flexion to 120

IV. 6 Weeks from Surgery

- Discontinue sling use
- Progress to full PROM and AAROM in all planes
- Wall alphabet with ball
- Active shoulder depressions with body weight
- UE gentle PNF (D1 and D2) AAROM with therapist
- Standing UE weight shifting
- Manual rhythmic stabilization at various angles
- Theraband Rows
- Biceps and triceps resistive strengthening
- Initiate AROM (flexion, scaption, ER 0 degrees of abduction)

V. 7 Weeks from Surgery

- Progress AROM
- Rotator cuff exercises at 0 degrees of abduction
- Quadruped stabilization
- Supine punches
- Refine shoulder mechanics

VI. 8 Weeks from Surgery

- Initiate UBE
- Full PROM
- Wall push ups
- Theraband exercises out of neutral
- UE PNF patterns
- Work on eccentric control of RTC muscles
- Light functional activities
- Chest press to neutral

VII. 10 Weeks from Surgery

• Shoulder extension exercises

VII. Weeks 12-20

- Include plyometric exercises (i.e. chest pass, overhead pass)
- Progress resistive exercises and shoulder intensive exercises
- Progress functional and sport specific activities